English	Tigrinya
What to expect after your COVID-19 vaccination	ድሕሪ ክታበት ኮቪድ-19 ምውሳድኩም እንታይ ትጽበዩ፡
Information for people who have just had a COVID-19 vaccination	ነቶም ክታበት ኮቪድ-19 ዝወሰዱ ሰባት ዝምልከት ሓበሬታ
Find out more at www.nidirect.gov.uk/covid-vaccine	ኣብዚ ዝስዕብ ነቝጣ ሙርበብ ሓበሬታ ተወሰኺ ሙብርሂ ድለዩ www.nidirect.gov.uk/covid-vaccine
Contents	ትሕዝቶታት
What are the side effects?	•
Can I catch COVID-19 from the vaccines?	• ካብቶም ክታበታት ሕማም ኮቪድ-19 ክሕዘኒ ይኽእልዶ፡
When can I go back to daily activities?	• ኣብ ንቡር ናይ ምዓልታዊ ንጥፈታተይ ሙዓስ ክምለስ እኽእል፡
Will the vaccine protect me?	•
What can I do after I have had the vaccine?	 ክታበት ምስ ተኸተብኩ እንታይ ክንብር እኽእል፡
Please read the product information leaflet for more details on your vaccine, including possible side effects, by searching Coronavirus Yellow Card.	ብኽብረትኩም ብዛዕባ ክታበታትኩም ዝምልከት፡ ነዚ ናይ ፍርያት ወረቐት ሓበሬታ ተወከሱ እንተላይ ንኸጋጥሙ ዝኽእሉ ጎድናዊ ሳዕቤናት፡ ናይ ኮሮናቫይረስ ብጫ ካርድ (Yellow Card) ፈትሹ፡፡
You can also report suspected side effects on the same website or by downloading the Yellow Card app.	ከምኡ'ውን ናይ ዝተጠርጠሩ ሳድናዊ ሳዕቤን ናብቲ ጦርበብ ሓበሬታ ወይ ነቲ ብጫ ካርድ <mark>ሞ</mark> መልከቲ ብምውራድ ጸብጻብ ክትህቡ ትኽእሉ።

https://coronavirus- yellowcard.mhra.gov.uk	https://coronavirus- yellowcard.mhra.gov.uk
People who are most at risk from the complications of coronavirus (COVID-19) are being offered the COVID-19 vaccination. This leaflet tells you what to expect after you have had your primary or booster vaccination.	ነቶም ኣብ ዝለዓለ ጸንም ናይ ኮሮናቫይረስ (ኮቪድ-19) ዝርከቡ ሰባት ክታበት ኮቪድ-19 እናተዋሀቦም ይርከብ። እዚ ንእሽቶ መሐበሪ ወረቐት እዚ ድሕሪ ምኽታብኩም እንታይ ክትጽበዩ ከምዘለኩም ዝንልጽ እዩ።
What are the side effects?	እቶም ሳድናውያን ሳዕቤናት እንታይ እዮም፥
Like all medicines, vaccines can cause side effects. Most of these are mild and short-term, and not everyone gets them. The very common side effects are the same and should still only last a day or two.	ከም ክሎም
Very common side effects in the first day or two include:	አብታ ቀዳሞይቲ ወይ ካልኣይቲ ሞዓልቲ ዘ <i>ጋ</i> ጥሙ ሳድናዊ ሳዕቤናት እነተም ዝስዕቡ እዮም፦
having a painful, heavy feeling and tenderness in the arm where you had your injection	ኣብታ ክታበት ዝወሰድክሙላ ኢድኩም ኣቐንዛውን ከቢድን ስምዒት ከምኡ'ውን ናይ ምንቅስቓስ ጸንምን ይህልወኩም።
feeling tiredheadache, aches and chills	 ስምዒት ድኻም ቃንዛ ርእሲ፡ ቅርጸትን ንብንብታን

You may also have flu like symptoms with episodes of shivering and shaking for a day or two. However, a high temperature* could also indicate that you have COVID-19 or another infection. You can rest and take the normal dose of paracetamol (follow the advice in the packaging) to help make you feel better.

ንሓደ ወይ ክልተ ሙዓልቲ ዝቅጵል ጉንፋዕ ዝሙስል ምልክታት ምንቅጥቃጥን ንብንብታን ዘጠቓለለ ስምዒት ምህላው። እንተኾነፃን፡ ልዑል ረስኒ ኣካላት* ምህላው ሕማም ኮቪድ-19 ወይ ካልእ ረኽሲ ከምዘለኩም የሙልክት (ነዚ ዝምልከት ምኽሪ ኣብ ንጽ 6 ተወከሱ)። ፃቡእ ዝኾነ ዓቐን ሙጠን ፈውሲ ቃንዛ ፓራሲታሞል የዕሪፍኩም ብምውሳድ (ነቲ ኣብ ናይ ሓበሬታ ወረቐት ተጠቒሱ ዘሎ ሙፃለጿ ብምኽታል) ጽቡቕ ንኽስምዓኩም ክትወስድዎ ትኽእሉ።

*If you are on chemotherapy and have a high temperature, contact your local cancer care unit.

An uncommon side effect is swollen glands in the armpit or neck on the same side as the arm where you had the vaccine. This can last for around 10 days, but if it lasts longer see your doctor. If you are due for a mammogram in the few weeks after the vaccine, then you should mention that when you attend.

ካልእ ዘይልሙድ ሳድናዊ ሳዕቤን ድማ ምሕባጥ ጵክታት ኣብ ትሽትሽ ወይ ኣብ ክሳድ ኣብ ሓደ ወንን ናይቲ ዝተኸተብካሉ ቦታ ከጋጥም ይኽእል። እዚ ኣስታት 10 ሙዓልታት ኣቢሉ ክጸንሕ ይኽእል፡ ካብኡ ንላዕሊ ምስ ዝጸንሕ ግን ናብ ሓኪምኩም ተራኣዩ። ንስኽን ቁሩብ ሰሙናት ድሕሪ ክታበት ናይ ማሞግራም (ኣብ ከባቢ ጡብ ዝግበር ሙርሙራ ራጂ) እትንብራሉ እዋን እንተበጲሑ፡ ኣብቲ ቦታ ክታበት ምስ ሙጻእክን ንሙዓስ ከምዝኾነ ቆጸራኽን ሓብራ።

What should I do if I am concerned about my symptoms?	ብዛዕባ እቶም ዝተርእዩኒ ምልክታት ሕማም ዘሰኩፉኒ ምስ ዝኾኑ እንታይ ክንብር ኣሎኒ፡
These symptoms normally last less than a week. If your symptoms seem to get worse or if you are concerned, call your GP or Out of Hours service. If you do seek advice from a doctor or nurse, make sure you tell them about your vaccination (show them the vaccination card – keep your card safe) so that they can assess you properly. You can also report suspected side effects of vaccines and medicines through the Yellow Card scheme. You can do this online by searching Coronavirus Yellow Card or by downloading the Yellow Card app.	እተም ምልክታት ሕጣም እዚኦም ካብ ሰሙን ዝውሕድ ግዜ ዝጸንሑ እዮም። እቶም ዝረኣይኹም ምልክታት እናበርትዑ ዝኾዱ ምስ ዝሙስልኹም ወይ ዝኾነ ስክፍታ ምስ ዝህልወኩም፡ ናብ ሓፈሻዊ ሓኪምኩም ወይ ናብቶም ካብ ሰዓታት ስራሕ ወጻኢ አንልግሎት ዝህቡ ኣላካት ደዉሉ። ካብ ሓኪም ወይ ካብ ኣላዪ ሕሙማት ምኽሪ ምስ ዘድልየኩም፡ ብዝዕባ ዝወሰድኩሞ ክታበት ከምዝነንርኩሞ ኣረጋግጹ (ንቲ ካርድ ክታበት ድማ ኣርእይዎም) ስለዚ ንሳቶም ብግቡእ ክምርምሩኹም ይኽእሉ። ብተወሳኺ ነቶም እትጥርጥርዎም ንድናዊ ሳዕቤናት ክታበት ከምኡውን መድሃኒታት በቲ ብጫ መሐበሪ ካርድ ኣቢልኩም ጸብጻብ ከተቅርቡ ትኽእሉ ኢኹም። ብተወሳኺ ንስኹም ንቲ ናይ ኮሮናቫይረስ ብጫ ካርድ ኣብ ቀጥታዊ መስመር ኢንተርነት ብምድላይ ወይ ነታ ናይ ኮሮናቫይረስ ብጫ ካርድ ብምጽዓን ክትንብርዎ ትኽእሉ፡ (ንዚ ብዝምልከት ብኽብረትኩም ኣብ ንጽ 2 ተመልከቱ)
https://coronavirus- yellowcard.mhra.gov.uk	https://coronavirus- yellowcard.mhra.gov.uk
Are there other more serious side effects?	ካልኦት ኣዝዮም ብርቱዓት <i>ጎ</i> ድናዊ ሳዕቤናት ኣለዉዶ፤
There have been reports of an extremely rare condition involving	ድሕሪ AstraZeneca ከምኡውን Janssen ዝተባህሉ ክታበታት ምኽታብ ኣብ ኣዝዩ

blood clots and unusual bleeding ዉሑድ *ኣጋ*ጣሚታት ተርእዮታት ምር*ጋ*እ after the AstraZeneca and Janssen vaccines. Because of the high risk of complications and death from ልዑል ሓደ*ጋ*ን ሕልኽላኻትን፡ ሞትን፡ ትካል COVID-19, the Medicines and Healthcare products Regulatory Agency (MHRA), the World Health MHRA፡ ትካል ጥዕና ዓለም፡ ትካል Organisation and the European <u>ምድሃኒ</u>ታት አውሮጳን ሚዛኑ ዝሓለወ Medicines Agency have concluded ክታበት ምህላዉ ኣዝዩ ከም ዘድሊ that the balance is very much in ደምዲሞም ኣለዉ። favour of vaccination. If you experience any of the ንስዥም ዝኾነ ካብዞም ዝስዕቡ following from more than 4 days ምልክታት ኣብ ውሽጢ 28 ማዓልታት ን and within 28 days after vaccination you should seek ምስ ዘ*ጋ*ጥሞኩም ካብ ሓፈሻዊ medical advice urgently from **ሓኪምኩም ወይ ካብ ክፍሊ ሀ**ጹጽ your GP or Emergency ረዲኤት ሕክምና ብዝቐልጠፈ ግዜ Department. ሕክምናዊ ሓንዝ ክትደልዩ ኣለኩም። • a new, severe headache • ሓድሽ፡ ብርቱዕ ቃንዛ ርእሲ ብልሙድ which is not helped by usual ፈውሲ ቃንዛ ዘይሓዊ ወይ እናበርት0 painkillers or is getting worse ዝ ኸ ይድ • an unusual headache which seems worse when lying ርእሲ ወይ ተዓዕጺጽኩም ኣብ down or bending over or may እትድቕስሉ *ግ*ዜ ዝብርትዕ ምስ be accompanied by ዝሞስል ወይ ምናልባት ብኻልእ ምልክት ዝተሰነየ ምስ ዝኸውን blurred vision, nausea – ዘይንጹር ምርኣይ፡ ዕግርግር and vomiting ምባልን ተምላስን፡ difficulty with your – ሽൗር ኣብ ኣዘራርባ፡ speech

 weakness, drowsiness or seizures 	– ድኽጦት፡ ትኽስታ ወይ ምንፍርፋር
 new, unexplained pinprick bruising or bleeding 	 ሓድሽ፡ ክግለጽ ዘይክ ል ል ሕጻጽ ወይ
 shortness of breath, chest pain, leg swelling or persistent abdominal pain 	• ሕጽረት እስትንፋስ፡ ቃንዛ ኣፍልቢ፡ ሕበጥ
Worldwide, rare cases of inflammation of the heart called myocarditis or pericarditis have been reported after Pfizer and Moderna COVID-19 vaccines.	ብደረጃ ዓለም፡ ኣዝዮም ውሑዳት ኣጋጣሚታት ነድሪ ጭዋዳታት ልቢ (myocarditis) ወይ ነድሪ ሽፋን ልቢ (pericarditis) ድሕሪ Pfizer ከምኡውን Moderna ዝተባህሉ ክታበታት ናይ ኮቪድ-19 ምኽታብ ኣጋጢሞም።
These cases have been seen mostly in younger men within several days after vaccination. Most of these people recovered and felt better following rest and simple treatments.	እዚ ተርእዮታት እዚ ኣውራ ኣብቶም ንኣሽቱ
You should seek medical advice urgently if you experience:	እዞም ዝስዕቡ ምልክታት ምስ ዝሀልውዥም ቅልጡፍ ሕክምናዊ ሓንዝ ክትደልዩ ኣለኩም፦
chest pain	• ቃንዛ ኣፍልቢ፡
shortness of breath	• ሕጽረት እስትንፋስ፡

 feelings of having a fastbeating, fluttering, or pounding heart

ስምዒት ቅልጡፍ ሀርመት ልቢ፡
 ድማድግታ ሀርመት ልቢ፡ ወይ ዝስማዕ
 ድምጺ ሀርመት ልቢ

Can I catch COVID-19 from the vaccines?

ካብ ክታበት ካቪድ-19 ሕማም ክሕዘኒ ይኽእልድዩ፡

You cannot catch COVID-19 from the vaccine but it is possible to have caught COVID-19 and not realise you have the symptoms until after your vaccination appointment. ካብቲ ክታበት ሕማም ኮቪድ-19 ክሕዘኩም ኣይክእልን እዩ፡ እንተኾነ ግን ከይተረድኣኩም ብኻልእ ሞኽንያት ሕማም ኮቪድ-19 ተታሒዝኩም ክትኮኑ ትኽእሉ እቲ ምልክታት ድማ ድሕሪ ቆጸራ ሞዓልቲ ክታበት ክርአ ይኽእል።

Although a fever can occur within a day or two of vaccination, if you have any other COVID-19 symptoms or your fever lasts longer, stay at home and arrange to have a test.

ዋላእሷ ኣብ ውሽጢ ሓደ ሙዓልቲ ወይ ክልተ ሙዓልታት ኣብ ግዜ ክታበት ረስኒ ናይ ኣካላት ዝርአ እንተኾነ፡ ዝኾነ ካልእ ምልክታት ሕማም ኮቪድ-19 ምስ ዝሀልወኩም ወይ ረስኒ ኣካላትኩም ንዝነውሐ ግዜ ምስ ዝጸንሕ፡ ኣብ ንዛዥም ጽንሑ ወይ ሙርሙራ ክግበረልኩም ቆጸራ ሓዙ።

When can I go back to daily activities?

ናብ ንቡር **ሞዓልታዊ ንጥፈታተይ ምዓስ** ክምለስ እኽእል፤

You should be able to resume activities that are normal for you as long as you feel well. If your arm is particularly sore, you may find heavy lifting difficult. If you feel unwell or very tired you should rest and avoid operating machinery or driving.

ጽቡቕ ክሳዕ ዝተሰማዓኩም ኣብ ዝኾነ ግዜ ንቡር ንጥፈታትኩም ክትጅሙሩ ትኽእሉ። ብፍላይ ኢድኩም እንተቖሲሉ፡ ዝኾነ ክብደት ኣብ ምልዓል ጸንም ከጋጥሙኩም ይኽእል እዩ። ጽቡቕ ምስ ዘይስሙዓኩም ወይ ኣዚኹም ምስ እትደኽሙ ከተዕርፉን ዝኾነ ማሽነሪ ከይትሕዙን ምምራሕ ሙኪና ከተወግዱን ኣለኩም።

Will the vaccine protect me?	እቲ ክታበት ካብ ሕማም ክከላኸለለይ ድዩ፡
The COVID-19 vaccines that you have had have been shown to reduce the chance of you suffering from COVID-19 disease.	እቲ ዝወሰድኩሞ ክታበት ናይ ኮቪድ-19፡ ተኽእሎ ስቓይ ናይ ሕማም ኮቪድ-19 ክንክየልኩም ይኽእል።
It may take a few weeks from the first dose for your body to build up protection. Your body should respond more quickly (after a few days) after any additional doses. Like all medicines, no vaccine is completely effective, so you should continue to take recommended precautions to avoid infection. Some people may still get COVID-19 despite having a vaccination, but this should be less severe.	ኣካላትኩም ካብቲ ክታበት ናይ ተጻዋርነት ብቕዓት ንኸጥሪ ዉሑዳት ሰሙናት ከድልዮ ይኽእል። ኣካላትኩም ኣብ ዝቐልጠፈ ግዜ (ድሕሪ ቁሩብ ሙዓልታት) ተወሳኺ ሙጠን ክታበት ምስተንበረልኩም ግብረሙልሲ ክህብ ኣለዎ። ከም ኩሎም ሙድሃኒታት፡ ዝኾን ክታበትውን ሙሉእ ብምሉእ ኣድማዒ ኣይከውንን፡ ስለዚ ረኽሲ ንምክልኻል ነቶም ተዋሂቦም ዘለዉ ጥንቃቐታት ብቐጻሊ ምክትታል ኣንዳሲ እዩ። ንለ ሰባት ተኸቲቦም እንከለዉ ሕማም ኮቪድ-19 ኽሕዛም ይኽእል እዩ፡ እንተኾነግን እቲ ሕማም ከቢድ ኣይከውንን።
What can I do after I have had the vaccine?	ክታበት ምስ ወሰድኩ እንታይ ክንብር ይ ማ በኣኒ፣
The vaccine cannot give you COVID-19 infection, and a full course will reduce your chance of becoming seriously ill. It will help to reduce the risk of you passing on the virus. So it is important to continue to follow current national guidance.	ክታበት ናይ ኮቪድ-19 ረኽሲ ከስዕበልኩም ኣይክእልን እዩ፡ ከምኡውን ምሉእ ክታበት ምውሳድ ብርቱ ሕማም ናይ ምሓዝ ተኽእሎ ይንክየልኩም። ንስዥም ክሳዕ ክንደይ ቫይረስ ከተመሓላልፉ ተኽእሎ ከምዘለኩም ግን እንፈልጦ ነገር የለን። ስለዚ ነቶም ህሉዋት መምርሒታት ምኽታል ኣንዳሲ እዩ።
To protect yourself and your family, friends and colleagues you must still:	ንንዛ እር እስ ሹም ከም ኡውን ስድራቤትኩም፡ ኣዕርኽት ሹም ከምኡውን መሳራሕት ሹም ንምክልኻል

	ሕጇውን ነዞም ዝስዕቡ ሞምርሒታት
	ክትክተሉ ኣለኩም፦
wear a face covering where	•
advised	ማ በሩ
practise social distancing	• ማሕበራዊ ርሕቀት ተጠቐሙ
wash your hands carefully	•
and frequently	ተሓጸቡ
open windows to let fresh air	• ጵሩይ ኣየር ንኽህሉ ሙሳኹቲ
in	ምኽፋት
follow the current guidance at	• ነዞም ሕጇ ዘለዉ
www.gov.uk/coronavirus	ዝስዕብ ነቝጣ
	ምክትታል <u>www.gov.uk/coronavirus</u>
Vaccination, helping to protect	ክታበት ምውሳድ ነቶም ኣዝዮም
those most vulnerable.	ተጠቃዕቲ ሰባት ንምክልኻል ይሕ <i>ግ</i> ዝ፡፡
If you need more information on	ብዛዕባ ክታበት ኮቪድ-19 ተወሳኺ ሓበሬታ
the COVID-19 vaccination please	ምስ ዘድልየኩም ብኽብረትኩም ኣብዚ
visit: <u>www.nidirect.gov.uk/covid-</u> vaccine	ዝስዕብ ነቝጣ
VACCINE	www.nidirect.gov.uk/covid-vaccine
Information correct at time of	ኣብ ማዜ ሕትጮት ዝተኣረጮ ሓበሬታ ነቲ ናይ
publication. For the latest version of this leaflet and alternative	<u>መወዳ</u> እታ ሕትመት ናይዛ ናይ ሓበሬታ
formats visit the PHA website	ወረቐት ከምኡውን ናይ ኣማራጺ ቅጥዕታት
www.publichealth.hscni.net	ዝምልከት ኣብዚ ነቝጣ
	ናይ ክፍሊ ህዝባዊ ሓለዋ ጥዕና ብጽሑ
	www.publichealth.hscni.net
Published September 2021 by the Public Health Agency.	<u>ኣጀ</u> ንስያ ህዝባዊ ጥዕና
,	

Tigrinya translation of What to expect after your COVID vaccination

- © Crown copyright 2021. This information was originally developed by the UK Health Security Agency and is used under the Open Government Licence v3.0